



Oxford Cambridge and RSA

A Level Mathematics A

H240/03 Pure Mathematics and Mechanics

Practice Paper – Set 1

Time allowed: 2 hours

You must have:

- Printed Answer Booklet

You may use:

- a scientific or graphical calculator

INSTRUCTIONS

- Use black ink. HB pencil may be used for graphs and diagrams only.
- Complete the boxes provided on the Printed Answer Booklet with your name, centre number and candidate number.
- Answer **all** the questions.
- **Write your answer to each question in the space provided in the Printed Answer Booklet.** If additional space is required, you should use the lined page(s) at the end of the Printed Answer Booklet. The question number(s) must be clearly shown.
- Do **not** write in the barcodes.
- You are permitted to use a scientific or graphical calculator in this paper.
- Final answers should be given to a degree of accuracy appropriate to the context.
- The acceleration due to gravity is denoted by $g \text{ m s}^{-2}$. Unless otherwise instructed, when a numerical value is needed, use $g = 9.8$.

INFORMATION

- The total mark for this paper is **100**.
- The marks for each question are shown in brackets [].
- **You are reminded of the need for clear presentation in your answers.**
- The Printed Answer Booklet consists of **16** pages. The Question Paper consists of **8** pages.

Formulae
A Level Mathematics A (H240)

Arithmetic series

$$S_n = \frac{1}{2}n(a+l) = \frac{1}{2}n\{2a+(n-1)d\}$$

Geometric series

$$S_n = \frac{a(1-r^n)}{1-r}$$

$$S_\infty = \frac{a}{1-r} \quad \text{for } |r| < 1$$

Binomial series

$$(a+b)^n = a^n + {}^nC_1 a^{n-1}b + {}^nC_2 a^{n-2}b^2 + \dots + {}^nC_r a^{n-r}b^r + \dots + b^n \quad (n \in \mathbb{N})$$

$$\text{where } {}^nC_r = {}_nC_r = \binom{n}{r} = \frac{n!}{r!(n-r)!}$$

$$(1+x)^n = 1 + nx + \frac{n(n-1)}{2!}x^2 + \dots + \frac{n(n-1)\dots(n-r+1)}{r!}x^r + \dots \quad (|x| < 1, n \in \mathbb{R})$$

Differentiation

$f(x)$	$f'(x)$
$\tan kx$	$k \sec^2 kx$
$\sec x$	$\sec x \tan x$
$\cot x$	$-\operatorname{cosec}^2 x$
$\operatorname{cosec} x$	$-\operatorname{cosec} x \cot x$

$$\text{Quotient rule } y = \frac{u}{v}, \quad \frac{dy}{dx} = \frac{v \frac{du}{dx} - u \frac{dv}{dx}}{v^2}$$

Differentiation from first principles

$$f'(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h}$$

Integration

$$\int \frac{f'(x)}{f(x)} dx = \ln|f(x)| + c$$

$$\int f'(x)(f(x))^n dx = \frac{1}{n+1}(f(x))^{n+1} + c$$

$$\text{Integration by parts } \int u \frac{dv}{dx} dx = uv - \int v \frac{du}{dx} dx$$

Small angle approximations

$\sin \theta \approx \theta$, $\cos \theta \approx 1 - \frac{1}{2}\theta^2$, $\tan \theta \approx \theta$ where θ is measured in radians

Trigonometric identities

$$\sin(A \pm B) = \sin A \cos B \pm \cos A \sin B$$

$$\cos(A \pm B) = \cos A \cos B \mp \sin A \sin B$$

$$\tan(A \pm B) = \frac{\tan A \pm \tan B}{1 \mp \tan A \tan B} \quad \left(A \pm B \neq \left(k + \frac{1}{2}\right)\pi\right)$$

Numerical methods

Trapezium rule: $\int_a^b y \, dx \approx \frac{1}{2}h\{(y_0 + y_n) + 2(y_1 + y_2 + \dots + y_{n-1})\}$, where $h = \frac{b-a}{n}$

The Newton-Raphson iteration for solving $f(x) = 0$: $x_{n+1} = x_n - \frac{f(x_n)}{f'(x_n)}$

Probability

$$P(A \cup B) = P(A) + P(B) - P(A \cap B)$$

$$P(A \cap B) = P(A)P(B|A) = P(B)P(A|B) \quad \text{or} \quad P(A|B) = \frac{P(A \cap B)}{P(B)}$$

Standard deviation

$$\sqrt{\frac{\sum(x-\bar{x})^2}{n}} = \sqrt{\frac{\sum x^2}{n} - \bar{x}^2} \quad \text{or} \quad \sqrt{\frac{\sum f(x-\bar{x})^2}{\sum f}} = \sqrt{\frac{\sum fx^2}{\sum f} - \bar{x}^2}$$

The binomial distribution

If $X \sim B(n, p)$ then $P(X = x) = \binom{n}{x} p^x (1-p)^{n-x}$, Mean of X is np , Variance of X is $np(1-p)$

Hypothesis test for the mean of a normal distribution

If $X \sim N(\mu, \sigma^2)$ then $\bar{X} \sim N\left(\mu, \frac{\sigma^2}{n}\right)$ and $\frac{\bar{X} - \mu}{\sigma/\sqrt{n}} \sim N(0, 1)$

Percentage points of the normal distribution

If Z has a normal distribution with mean 0 and variance 1 then, for each value of p , the table gives the value of z such that $P(Z \leq z) = p$.

p	0.75	0.90	0.95	0.975	0.99	0.995	0.9975	0.999	0.9995
z	0.674	1.282	1.645	1.960	2.326	2.576	2.807	3.090	3.291

Kinematics

Motion in a straight line

$$v = u + at$$

$$s = ut + \frac{1}{2}at^2$$

$$s = \frac{1}{2}(u + v)t$$

$$v^2 = u^2 + 2as$$

$$s = vt - \frac{1}{2}at^2$$

Motion in two dimensions

$$\mathbf{v} = \mathbf{u} + \mathbf{a}t$$

$$\mathbf{s} = \mathbf{u}t + \frac{1}{2}\mathbf{a}t^2$$

$$\mathbf{s} = \frac{1}{2}(\mathbf{u} + \mathbf{v})t$$

$$\mathbf{s} = \mathbf{v}t - \frac{1}{2}\mathbf{a}t^2$$

Section A: Pure Mathematics

Answer **all** the questions**1 In this question you must show detailed reasoning.**

Find the gradient of the curve $y = 3 \cos 2x$ at the point where $x = \frac{1}{8}\pi$. [4]


2 (i) Express $4 \cos \theta + 3 \sin \theta$ in the form $R \cos(\theta - \alpha)$, where $R > 0$ and $0^\circ < \alpha < 90^\circ$. [3]

The temperature $\theta^\circ\text{C}$ of a building at time t hours after midday is modelled using the equation

$$\theta = 20 + 4 \cos(15t)^\circ + 3 \sin(15t)^\circ, \text{ for } 0 \leq t < 24.$$

(ii) Find the minimum temperature of the building as given by this model. [1]

(iii) Find also the time of day when this minimum temperature occurs. [3]

3

The diagram shows triangle ABC , in which angle $A = \theta$ radians, angle $B = \frac{3}{4}\pi$ radians and $AB = 1$ unit.

(i) Use the sine rule to show that $AC = \frac{1}{\cos \theta - \sin \theta}$. [3]

(ii) Given that θ is a small angle, use the result in part **(i)** to show that

$$AC \approx 1 + p\theta + q\theta^2,$$

where p and q are constants to be determined. [4]

4 In this question you must show detailed reasoning.

It is given that the geometric series

$$1 + \frac{5}{3x-4} + \left(\frac{5}{3x-4}\right)^2 + \left(\frac{5}{3x-4}\right)^3 + \dots$$

is convergent.

(i) Find the set of possible values of x , giving your answer in set notation. [5]

(ii) Given that the sum to infinity of the series is $\frac{2}{3}$, find the value of x . [3]

- 5 (i) By sketching the graphs of $y = \frac{5}{x^2}$ and $y = |2 - 4x|$ on a single diagram, show that the equation

$$\frac{5}{x^2} = |2 - 4x| \quad (\text{A})$$

has exactly two real roots. [3]

- (ii) Show that the positive root α of equation (A) satisfies the equation $f(x) = 0$, where

$$f(x) = 4x^3 - 2x^2 - 5. \quad [1]$$

- (iii) Hence show that a Newton-Raphson iterative formula for finding α can be written in the form


$$x_{n+1} = \frac{8x_n^3 - 2x_n^2 + 5}{12x_n^2 - 4x_n}. \quad [3]$$

- (iv) Use this iterative formula, with initial value $x_1 = 1$, to find the value of α correct to 3 decimal places. Show the result of each iteration. [3]

A student claims that the iterative formula from part (iii) can be used to find the negative root of equation (A) provided that a suitable initial value is chosen.

- (v) Explain why the student's claim is incorrect. [1]

- 6 (i) Show that the two non-stationary points of inflection on the curve $y = \ln(1 + 4x^2)$ are at $x = \pm \frac{1}{2}$. [6]



The diagram shows the curve $y = \ln(1 + 4x^2)$. The shaded region is bounded by the curve and a line parallel to the x -axis which meets the curve where $x = \frac{1}{2}$ and $x = -\frac{1}{2}$.

- (ii) Show that the area of the shaded region is given by

$$\int_0^{\ln 2} \sqrt{e^y - 1} \, dy. \quad [3]$$

- (iii) Show that the substitution $e^y = \sec^2 \theta$ transforms the integral in part (ii) to $\int_0^{\frac{1}{4}\pi} 2 \tan^2 \theta \, d\theta$. [2]

- (iv) Hence find the exact area of the shaded region. [3]

Section B: Mechanics
Answer **all** the questions

- 7 A monorail train travels along a straight horizontal track from station A to station B . The train accelerates uniformly from rest at A to a maximum speed of 30 m s^{-1} then travels at this speed for 90 seconds before slowing down uniformly to come to rest at B . The acceleration of the train is $a \text{ m s}^{-2}$, the deceleration is $2a \text{ m s}^{-2}$ and the time for the whole journey is 3 minutes.
- (i) Sketch the velocity-time graph for the journey. [2]
- (ii) Calculate the distance between the two stations. [2]
- (iii) Calculate the value of a . [3]
- 8 A particle P of mass 3 kg moves under the action of a force $\begin{pmatrix} 9 \\ -3 \end{pmatrix} \text{ N}$. Initially P has velocity $\begin{pmatrix} 1 \\ -2 \end{pmatrix} \text{ m s}^{-1}$ and is at the point with position vector $\begin{pmatrix} 2 \\ 3 \end{pmatrix} \text{ m}$. At time t seconds later, P has velocity $\mathbf{v} \text{ m s}^{-1}$.
- (i) Express \mathbf{v} in terms of t . [2]
- (ii) Find the value of t when the speed of P reaches 5 m s^{-1} . [3]
- (iii) Find the position vector of P when $t = 2$. [2]
- 9 A boy kicks a ball from a point O on horizontal ground. The ball first hits the ground at a distance of 60 m from O and the time of flight is 4 seconds. This motion of the ball is modelled as that of a particle moving freely under gravity.
- (i) Find the horizontal and vertical components of the initial velocity of the ball. [3]
- The ball just clears a vertical post, of height h m, at a horizontal distance of 15 m from O .
- (ii) Show that $h = 14.7$. [2]
- (iii) Find the speed of the ball as it passes over the post. [4]
- Measurements show that the speed of the ball as it passes over the post is in fact not equal to the value found in part (iii).
- (iv) State a deficiency of the model that might account for this. [1]
- (v) Explain whether an improved model would require a larger or smaller initial speed for the ball. [1]

- 10** A particle P of weight W lies on the surface of a rough plane which is inclined at an angle α to the horizontal, where $\tan \alpha = \frac{4}{3}$. The coefficient of friction between the particle and the plane is $\frac{1}{2}$. A horizontal force of magnitude H is applied to P . This force acts in the vertical plane through a line of greatest slope. It is given that H is the greatest value for which P remains in equilibrium.


(i) Indicate on a diagram the forces acting on P . [1]

(ii) Show that $H = \frac{11}{2}W$. [5]

The horizontal force acting on P is now removed.

(iii) Find the acceleration of P in terms of g . [4]

11



A thin light rod AD has length $3a$. The end A is in contact with a smooth vertical wall which is perpendicular to the vertical plane containing the rod. The rod carries a load of weight W at the end D . The rod is held in equilibrium by two fixed smooth pegs B and C , where $AB = BC = CD = a$. The rod passes under peg B and over peg C , and makes an angle θ with the horizontal (see diagram).

(i) (a) Show that the normal contact force at C may be expressed as $W\left(\frac{3\cos^2\theta - 1}{\cos\theta}\right)$. [5]

(b) Find the normal contact force at B in terms of W and θ . [1]

(ii) Hence show that the value of θ is at most 35.3° , correct to 3 significant figures. [2]

(iii) Show that it is not possible for the magnitude of the reaction at A to equal the magnitude of the reaction at C . [6]

END OF QUESTION PAPER

OCR

Oxford Cambridge and RSA

Copyright Information

OCR is committed to seeking permission to reproduce all third-party content that it uses in its assessment materials. OCR has attempted to identify and contact all copyright holders whose work is used in this paper. To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced in the OCR Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download from our public website (www.ocr.org.uk) after the live examination series.

If OCR has unwittingly failed to correctly acknowledge or clear any third-party content in this assessment material, OCR will be happy to correct its mistake at the earliest possible opportunity.

For queries or further information please contact the Copyright Team, First Floor, 9 Hills Road, Cambridge CB2 1GE.

OCR is part of the Cambridge Assessment Group; Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.